

FACTOR CONGRUENCES IN SEMILATTICES

PEDRO SÁNCHEZ TERRAF

ABSTRACT. We characterize factor congruences in semilattices by using generalized notions of order ideal and of direct sum of ideals. When the semilattice has a minimum (maximum) element, these generalized ideals turn into ordinary (dual) ideals.

1. INTRODUCTION

In this paper we characterize direct factors in semilattices by using generalized notions of order ideal and of direct sum of ideals. When the semilattice has a minimum (maximum) element, these generalized ideals turn into ordinary (dual) ideals.

The main feature of this characterization is that it is completely analog to similar definitions in the realm of classical algebra, as in ring theory. By using the partial meet operation in a join-semilattice, we devise a definition with the same (quantifier) complexity as the reader may encounter in classical algebra, and we obtain an equational relation between each element and its direct summands in a given decomposition.

By the end of the paper we apply the characterization to the bounded case and show that the each of the axioms is necessary.

2. GENERALIZED DIRECT SUMS

The key idea in our characterization stems from the fact that in a direct product of join-semilattices there must exist “non-trivial” meets satisfying certain modularity and absorption laws with respect to join. And conversely, the existence of meets in a direct product implies the existence of them in each factor. We state without proof the latter property:

Claim 1. *For every $\vee$ -semilattices C, D and elements $c, e \in C$ and $d, f \in D$, if $(c, d) \wedge (e, f)$ exists in $C \times D$, then $c \wedge e$ and $d \wedge f$ exist (and conversely).*

In the following, we will write formulas in the extended language $\{\vee, \wedge\}$. The subformula “ $x \wedge y = z$ ” will interpreted as “ $x \wedge y$ exists and equals z ”:

$$\exists w : w \leq x, y \ \& \ (\forall u : u \leq x, y \rightarrow u \leq w) \ \& \ w = z,$$

unless explicitly indicated, and every occurrence of $\wedge$ will be reduced by means of this procedure. It is easy to see that the associative law holds for the partial $\wedge$ operation in every $\vee$ -semilattice, and we will apply it without any further mention.

Supported by Conicet.

Keywords: semilattice, direct factor, factor congruence, generalized direct sum, generalized ideal.

MSC 2000: 06A12, 20M10.

Heretofore, A will be a $\vee$ -semilattice and $c \in A$ arbitrary but fixed. Let $\phi_c(x_1, x_2, x)$ be the conjunction of the following formulas:

dist $x = (x \vee x_1) \wedge (x \vee x_2)$.
p1 $x_1 = (x \vee x_1) \wedge (c \vee x_1)$.
p2 $x_2 = (x \vee x_2) \wedge (c \vee x_2)$.
join $x_1 \vee x_2 = x \vee c$.

Definition 2. Assume I_1, I_2 are subsemilattices of A . We will say that A is the c -direct sum of I_1 and I_2 (notation: $A = I_1 \oplus_c I_2$) if and only if the following hold:

Mod1 For all $x, y \in A$, $x_1 \in I_1$ and $x_2 \in I_2$, if $x \vee c \geq x_1 \vee x_2$ then

$$((x \vee x_1) \wedge (x \vee x_2)) \vee y = (x \vee y \vee x_1) \wedge (x \vee y \vee x_2).$$

Mod2 For all $x, y \in A$, $x_1 \in I_1$ and $x_2 \in I_2$, if $x \leq x_1 \vee x_2$ then

$$((x \vee x_i) \wedge (c \vee x_i)) \vee y = (x \vee y \vee x_i) \wedge (c \vee y \vee x_i)$$

for $i = 1, 2$.

Abs For all $x_1, y_1 \in I_1$ and $z_2 \in I_2$, if any of the following meets exists, the other one also does, and we have : $x_1 \wedge (y_1 \vee z_2) = x_1 \wedge (y_1 \vee c)$ (and interchanging I_1 and I_2).

exi $\forall x_1 \in I_1, x_2 \in I_2 \exists x \in A : \phi_c(x_1, x_2, x)$.

onto $\forall x \in A \exists x_1 \in I_1, x_2 \in I_2 : \phi_c(x_1, x_2, x)$.

In order to make notation lighter, we will drop the subindex “ c ”. Let us notice that **exi** implies:

ori $\forall x_1 \in I_1, x_2 \in I_2 (x_1 \vee x_2 \geq c)$.

Lemma 3. Assume $A = I_1 \oplus I_2$. Then $\phi(x_1, x_2, x)$ defines an isomorphism $(x_1, x_2) \xrightarrow{\phi} x$ between $I_1 \times I_2$ and A .

Proof. First we'll see that the function $(x_1, x_2) \xrightarrow{\phi} x$ is well defined. Let us suppose $\phi(x_1, x_2, x)$ and $\phi(x_1, x_2, z)$ (there is at least one possible image by **exi**). We operate as follows:

$$\begin{aligned} z &= (z \vee x_1) \wedge (z \vee x_2) && \text{by } \mathbf{dist} \text{ for } z \\ &= [z \vee ((x \vee x_1) \wedge (c \vee x_1))] \wedge \\ &\quad \wedge [z \vee ((x \vee x_2) \wedge (c \vee x_2))] && \text{by } \mathbf{p1} \text{ and } \mathbf{p2}. \end{aligned}$$

By **join** for x we may apply **Mod2**:

$$= [(z \vee x \vee x_1) \wedge (\underline{c} \vee z \vee x_1)] \wedge [(z \vee x \vee x_2) \wedge (\underline{c} \vee z \vee x_2)]$$

and by **join** for z ,

$$= [(x \vee z \vee x_1) \wedge (x_1 \vee x_2)] \wedge [(x \vee z \vee x_2) \wedge (x_1 \vee x_2)]$$

This last term is symmetric in x and z , hence we obtain $x = z$. The function thus defined by ϕ is surjective by **onto**; let us now prove that it is 1-1. If $\phi(x_1, x_2, x)$ and $\phi(y_1, y_2, x)$, we have:

$$\begin{aligned} x_1 &= x_1 \wedge (x_1 \vee x_2) \\ &= x_1 \wedge (x \vee c) && \text{by } \mathbf{join} \\ &= x_1 \wedge (y_1 \vee y_2) && \text{by } \mathbf{join} \text{ again} \\ &= x_1 \wedge (y_1 \vee c) && \text{by } \mathbf{Abs} \end{aligned}$$

hence $x_1 \leq y_1 \vee c$. Symmetrically, $y_1 \leq x_1 \vee c$ and in conclusion

$$(1) \quad x_1 \vee c = y_1 \vee c.$$

On the other hand,

$$\begin{aligned} x \vee y_1 &= ((x \vee x_1) \wedge (x \vee x_2)) \vee y_1 && \text{by } \mathbf{dist} \\ &= (x \vee y_1 \vee x_1) \wedge (x \vee y_1 \vee x_2) && \text{by } \mathbf{Mod1} \\ &= (x \vee y_1 \vee x_1) \wedge (x \vee y_1 \vee x_2 \vee c) && \text{by } \mathbf{ori} \\ &= (x \vee y_1 \vee x_1) \wedge (x \vee c) && \text{since } x \vee c \geq x_1, y_1 \\ &= x \vee y_1 \vee x_1 && \text{by same argument.} \end{aligned}$$

Symmetrically,

$$x \vee x_1 = x \vee y_1 \vee x_1.$$

and hence $x \vee x_1 = x \vee y_1$. Collecting this with (1) and using **p₁** we have

$$x_1 = (x \vee x_1) \wedge (c \vee x_1) = (x \vee y_1) \wedge (c \vee y_1) = y_1.$$

By the same reasoning, $x_2 = y_2$.

Let us suppose that $\phi(x_1, x_2, x)$ and $\phi(z_1, z_2, z)$; since each of I_1, I_2 is a subsemilattice, we know that $x_j \vee z_j \in I_j$ for $j = 1, 2$. We will see that $\phi(x_1 \vee z_1, x_2 \vee z_2, x \vee z)$ (i.e., ϕ preserves $\vee$). The property **join** is immediate. We now prove **dist**:

$$\begin{aligned} (2) \quad x \vee z &= ((x \vee x_1) \wedge (x \vee x_2)) \vee z && \text{by } \mathbf{dist} \text{ for } x \\ (3) \quad &= (x \vee x_1 \vee z) \wedge (x \vee x_2 \vee z) && \text{by } \mathbf{Mod1} \\ (4) \quad &= (x \vee z \vee x_1 \vee z_1) \wedge (x \vee z \vee x_1 \vee z_2) \wedge \\ (5) \quad &\quad \wedge (x \vee z \vee x_2 \vee z_1) \wedge (x \vee z \vee x_2 \vee z_2) \end{aligned}$$

by **dist** for z and **Mod1**. Note that

$$\begin{aligned} x \vee z \vee x_1 \vee z_2 &= x \vee c \vee z \vee c \vee x_1 \vee z_2 && \text{by } \mathbf{ori}, x_1 \vee z_2 = x_1 \vee z_2 \vee c \\ &= x_1 \vee x_2 \vee z_1 \vee z_2 \vee x \vee z && \text{by } \mathbf{join} \text{ two times} \\ &\geq x \vee z \vee x_1 \vee z_1, \end{aligned}$$

hence we can eliminate two terms in equation (5) and we obtain

$$(x \vee z) = ((x \vee z) \vee (x_1 \vee z_1)) \wedge ((x \vee z) \vee (x_2 \vee z_2)).$$

We may obtain $\mathbf{p}_1$ and $\mathbf{p}_2$ similarly. We prove $\mathbf{p}_1$:

$$\begin{aligned} x_1 \vee z_1 &= ((x \vee x_1) \wedge (c \vee x_1)) \vee z_1 && \text{by } \mathbf{p}_1 \text{ for } x \\ &= (x \vee z_1 \vee x_1) \wedge (c \vee z_1 \vee x_1) && \text{by } \mathbf{Mod2} \end{aligned}$$

By $\mathbf{p}_1$ for z followed by $\mathbf{Mod2}$ in each term of the meet,

$$(6) \quad = (x \vee z \vee x_1 \vee z_1) \wedge (x \vee c \vee x_1 \vee z_1) \wedge (c \vee z \vee x_1 \vee z_1) \wedge (c \vee x_1 \vee z_1)$$

and this equals

$$((x \vee z) \vee (x_1 \vee z_1)) \wedge (c \vee (x_1 \vee z_1)).$$

since the last term in (6) is less than or equal to the previous ones. $\square$

Since $\phi(\cdot, \cdot, \cdot)$ defines an isomorphism (relative to I_1, I_2), there are canonical projections $\pi_j : A \rightarrow I_j$ with $j = 1, 2$ such that $\phi(\pi_1(x), \pi_2(x), x)$ holds.

Theorem 4. *The mappings*

$$\begin{aligned} (\theta, \delta) &\longmapsto (I_\delta, I_\theta) \\ (\ker \pi_1, \ker \pi_2) &\longleftarrow (I_1, I_2) \end{aligned}$$

where $I_\theta := \{a \in A : a \theta c\}$, are mutually inverse maps defined between the set of pairs of complementary factor congruences of A and the set of pairs of subsemilattices I_1, I_2 of A such that $A = I_1 \oplus_c I_2$.

Proof. We only prove that the left-to-right map is well defined, i.e. $I_\delta \oplus_c I_\theta = A$ for every pair of complementary factor congruences θ, δ . It's clear that I_δ, I_θ are subsemilattices of A , and we know that the mapping $a \mapsto (a/\theta, a/\delta)$ is an isomorphism between A and $A/\theta \times A/\delta$. Under this isomorphism, I_δ corresponds to $\{(a', c') : a' \in A/\theta\}$ and I_θ to $\{(c', a'') : a'' \in A/\delta\}$, where $c' = c/\theta$ and $c'' = c/\delta$. We will check the axioms for $I_\delta \oplus_c I_\theta = A$ in the isomorphic image $A/\theta \times A/\delta$.

In order to see **Abs**, let $x_1 = (x, c'')$, $y_1 = (y, c'')$ and $z_2 = (c', z)$ as in the hypothesis and suppose that $x_1 \wedge (y_1 \vee z_2)$ exists. That is to say,

$$(x, c'') \wedge [(y, c'') \vee (c', z)] = (x, c'') \wedge (y \vee c', c'' \vee z)$$

exists. By Claim 1, we know that $x \wedge (y \vee c')$ must exist in A_1 , and in conclusion,

$$\begin{aligned} (x, c'') \wedge [(y, c'') \vee (c', z)] &= (x \wedge (y \vee c'), c'' \wedge (c'' \vee z)) = \\ &= (x \wedge (y \vee c'), c'') = (x, c'') \wedge [(y, c'') \vee (c', c'')]. \end{aligned}$$

That is to say, the other meet exists and they are equal. The other half is analogous, and the verification of **Mod1** and **Mod2** is a straightforward matter.

Let $x = (x', x'')$, $x_1 = (x', c'')$ and $x_2 = (c', x'')$. The map $I_\delta \times I_\theta \rightarrow A/\theta \times A/\delta$ given by $(x_1, x_2) \mapsto x$ is a bijection. Hence, to check **exi** and **onto** we just have to check that $\phi_c(x_1, x_2, x)$ holds:

dist We have:

$$\begin{aligned}
 x &= (x', x'') \\
 &= (x' \vee x', x'' \vee c'') \wedge (x' \vee c', x'' \vee x'') \\
 &= [(x', x'') \vee (x', c'')] \wedge [(x', x'') \vee (c', x'')] \\
 &= (x \vee x_1) \wedge (x \vee x_2),
 \end{aligned}$$

which is what we were looking for.

p₁ We reason as follows

$$\begin{aligned}
 x_1 &= (x', c'') \\
 &= (x' \vee x', x'' \vee c'') \wedge (x' \vee c', c'' \vee c'') \\
 &= [(x', x'') \vee (x', c'')] \wedge [(c', c'') \vee (x', c'')] \\
 &= (x \vee x_1) \wedge (c \vee x_1),
 \end{aligned}$$

p₂ It's totally analogous.

join Obvious. □

Remark. Since semilattices have BFC (see [1]), it can be seen that the set of direct summands of A , $\{I \leq A : \exists J \leq A, I \oplus J = A\}$, is a Boolean algebra and $\theta \rightarrow I_\theta$ is a lattice isomorphism from factor congruences to direct summands.

3. SOME PARTICULAR CASES

In the case our semilattice A has a 0 or a 1 —minimum or maximum element, respectively— the characterization takes a much simpler form. By taking c to be minimum (maximum), it turns out that the subsemilattices I_1 and I_2 are order (dual) ideals.

Theorem 5. *Let A be a semilattice with 0. Then $A = I_1 \oplus_0 I_2$ if and only if $I_1, I_2 \leq A$ satisfy:*

Abs *For all $x_1, y_1 \in I_1$ and $z_2 \in I_2$, if any of the following meets exists, the other one also does, and we have: $x_1 \wedge (y_1 \vee z_2) = x_1 \wedge y_1$ (and interchanging I_1 and I_2).*

onto $I_1 \vee I_2 = A$.

Moreover, I_1 and I_2 are ideals of A .

Proof. Since $y_1 \vee 0 = y_1$, **Abs** is equivalent to its new form as stated and **Mod1**, **Mod2** are trivially true. The **join** part of ϕ_0 reduces to “ $x_1 \vee x_2 = x$ ” and from here we conclude that **dist**, **p₁** and **p₂** hold trivially. Hence **exi** simplifies to

$$\forall x_1 \in I_1, x_2 \in I_2 \exists x \in A : x_1 \vee x_2 = x,$$

which holds trivially, and **onto** reduces to:

$$\forall x \in A \exists x_1 \in I_1, x_2 \in I_2 : x_1 \vee x_2 = x.$$

and this is equivalent to $I_1 \vee I_2 = A$.

To see that both of I_1 and I_2 are ideals, we observe that by Theorem 4 each of them is the θ -block of 0 for some congruence θ , hence they are downward closed. $\square$

Theorem 6. *Let A be a semilattice with 1. Then $A = I_1 \oplus_1 I_2$ if and only if $I_1, I_2 \leq A$ satisfy:*

- exi'** $I_1 \vee I_2 = \{1\}$
- onto'** $I_1 \wedge I_2 = A$.
- Mod1** $\forall x_1 \in I_1, x_2 \in I_2 : \forall y (x_1 \wedge x_2) \vee y = (y \vee x_1) \wedge (y \vee x_2)$,

Moreover, I_1 and I_2 are dual ideals (viz. upward closed sets) of A .

Proof. We may eliminate **Abs** and **Mod2** since they're trivial when $c = 1$ (this can be seen by considering **ori** after Definition 2). Formulas **p₁** and **p₂** turn into

$$x_1 = x \vee x_1 \quad x_2 = x \vee x_2$$

which are equivalent to say $x_1, x_2 \geq x$. Using this, we reduce **dist** to:

$$x = x_1 \wedge x_2,$$

and then we may eliminate **p₁** and **p₂** in favor of this last formula. **join** is equivalent to $x_1 \vee x_2 = 1$, and hence **exi** reduces to the statement “for all $x_1 \in I_1, x_2 \in I_2$, $x_1 \vee x_2 = 1$ and $x_1 \wedge x_2$ exists”. We may state the first part as $I_1 \vee I_2 = \{1\}$, and may condense the second part with **onto** writing $I_1 \wedge I_2 = A$. It remains to take care of **Mod1**. The hypothesis $x \vee c \geq x_1 \vee x_2$ trivializes when $c = 1$, and if we take (without loss of generality) $x := x_1 \wedge x_2$, we obtain

$$\forall x_1 \in I_1, x_2 \in I_2 : \forall y (x_1 \wedge x_2) \vee y = (y \vee x_1) \wedge (y \vee x_2),$$

which was to be proved. $\square$

4. INDEPENDENCE AND NECESSITY OF THE AXIOMS

We check independence and necessity by providing semilattices with a distinguished element c and a pair of ideals I_1 and I_2 such that they satisfy every axiom minus one and it is not the case that $A = I_1 \oplus_c I_2$.

It is immediate that the minimal non-modular lattice N_5 provide two such counterexamples showing that **Abs** and **Mod1** are necessary and independent (see Figure 1 (a) and (b)). Note that **Abs** implies $I_1 \cap I_2 = \{c\}$, but this sole assumption does not suffices to prove uniqueness of representations: in Figure 1 (a) we have $\phi_c(x_1, z_2, 1)$ and $\phi_c(y_1, z_2, 1)$.

To see that **Mod2** is necessary and independent is almost as easy as with the other axioms. Consider the semilattice pictured in Figure 1 (c). First note that every element in I_2 is greater than or equal to every element of I_2 . From this observation it is immediate to see that **Mod1** and **Abs** hold and that $\phi_c(x, y, z)$ (where $x \in I_1$ and $y \in I_2$) may be




FIGURE 1. Counterexamples.

simplified as follows:

$$\begin{array}{ll}
 z = (z \vee x) \wedge (z \vee y) = z \vee x & \text{since } x \leq y \\
 x = (z \vee x) \wedge (c \vee x) = z \wedge c & \text{by the previous line and } c \geq x \\
 y = (z \vee y) \wedge (c \vee y) = (z \wedge y) \vee y & \text{(holds trivially)} \\
 x \vee y = z \vee c & \text{and hence } y = z \vee c
 \end{array}$$

Therefore, $\phi_c(x, y, z)$ is equivalent to $(x = z \wedge c) \& (y = z \vee c)$ and it is now evident that **exi** and **onto** also hold. Finally, consider the labeled elements x , x_1 , x_2 and y . We have $x \leq x_1 \vee x_2$, and if **Mod2** were true we should have:

$$\begin{aligned}
 ((x \vee x_1) \wedge (c \vee x_1)) \vee y &= (x \vee y \vee x_1) \wedge (c \vee y \vee x_1) \\
 (x \wedge c) \vee y &= x_2 \wedge c \\
 x_1 \vee y &= c \\
 y &= c,
 \end{aligned}$$

which is an absurdity.

The last two counterexamples show that **onto** and **exi** are both independent and necessary. In Figure 2 (a) there exist no pair $(x_1, x_2) \in I_1 \times I_2$ such that $\phi_c(x_1, x_2, x)$, and in Figure 2 (b) there exists no x such that $\phi_c(x_1, x_2, x)$.

REFERENCES

- [1] C. C. Chen: On the unique factorization property of algebras. Nanta Math. **10** (1977), 149–157.

CIEM — Facultad de Matemática, Astronomía y Física (Fa.M.A.F.)
 Universidad Nacional de Córdoba - Ciudad Universitaria
 Córdoba 5000. Argentina.
 sterraf@mate.uncor.edu




FIGURE 2. Counterexamples.